## POSITIVE TB SKIN TEST (TST) / IGRA REPORTING & MEDICATION ORDER FORM

Public Health Services www.hamilton.ca/tuberculosis Phone: 905-546-2063 Fax: 1-844-444-0295 Hamilton

Please complete and fax this form and chest x-ray report to 905-546-4078 within 7 days.						
Patient's Last Name, First Name Middle Name			Date of Birt	h	Gender	
Tallotto Last Hamo, First Hamo illiadio Hamo					☐ Female ☐ Male	
			(dd/mmn	n/yyyy)	☐ Transgender ☐ Other	
Address, City, Postal Code		Home Phone Numl	ber Cell Phon		e Number	
Born in Canada	☐ Yes - Province:		Country of	Birth	Date of Arrival	
☐ Yes - identify as: ☐ First Nation						
☐ Inuit ☐ Other Indigenous					(dd/mmm/yyyy)	
Reason for Test		Routine screening (includes volunteer, school, work)  Medical   Immigration			History of BCG	
	│	•			☐ Unknown ☐ No ☐ Yes - Year:	
TST	Date Planted:				mm induration	
	(dd/mmm/yyyy)	(dd/mmm/)	yyy)			
IGRA	(dd/mmm/yyyy)					
Positive TST: ✓ 10 mm or more is considered positive for most people						
✓ 5 mm or more may be considered positive in specific situations listed in the Canadian TB Standards, 8 <sup>th</sup> Ed., Chapter 4, Table 1						
Patients with positive TST/IGRA require: ✓ Symptom assessment and physical exam						
✓ Chest x-ray - Date: Fax report to public health						
Symptom Assessment						
□ Asymptomatic □ Symptomatic - Specify: □ cough □ fever □ night sweats □ fatigue □ other:						
If symptomatic or chest x-ray indicates TB disease: ✓ Instruct patient to isolate at home (provide masks)						
✓ Collect 3 sputum specimens at least 1 hour apart						
✓ Report immediately to public health at 905-546-2063						
Risk Factors for TB Disease Progression (check all that apply)						
□ No risk factors □ Receiving immunosuppressive drugs					drugs	
☐ HIV infection			☐ Biologics			
· · · · · · · · · · · · · · · · · · ·			Moderate to high dose steroids			
☐ Age when infected - under 5 years			☐ Cancer (lung, sarcoma, leukemia, lymphoma or			
☐ Silicosis	•	gastrointestinal)				
☐ Chronic renal fail	☐ Diab					
☐ Transplant recipient			nol use (3 or r		- 7	
☐ Fibronodular disease			☐ Tobacco cigarette use (1 or more packs/day)			
☐ Granuloma on chest x-ray ☐ Underweight (less than 90% ideal body weight)						
<b>Note:</b> Refer to The Online TST/IGRA Interpreter Tool The Online TST/IGRA Interpreter to assess risk for active TB disease.						
Health Education and Follow-Up (check all that apply)						
☐ Reviewed signs & symptoms of active TB and when to seek health care ☐ Referred to family physic					erred to family physician	
☐ TB information provided - available at www.hamilton.ca/tuberculo			is ☐ Treatment discussed			
☐ Treatment prescribed (refer to TPT Prescription Section)				☐ Treatment refused		
☐ Referred to TB Clinic (Phone: 905-522-1155, Ext. 34198 Fax: 905-525-5806)						
TB Preventive Treatment (TPT) Prescription (Refer to TPT Quick Reference)						
Medication	Prescrip	otion (oral daily)	Mitte	e (in month	s) Duration (in months)	
Rifampin		mg 🛘 other:	_mg	1	4	
Isoniazid	☐ 300mg ☐ othe			1	9	
Pyridoxine (vitamin	, ,	r:mg		1	9	
* Vitamin B6 is given with isoniazid to minimize the risk of neuropathy						
Health Care Provider Name:			) #:		Date:	
Signature: Address: Phone: Fax:						